

CONDENSED DESCRIPTION OF MAP UNITS

The complete description of map units and references is in the accompanying booklet.

SURFICIAL DEPOSITS

HUMAN-MADE DEPOSITS

- af Artificial fill (latest Holocene)

ALLUVIAL DEPOSITS

- Modern stream channel, flood plain, and low terrace deposits (Holocene and late Pleistocene)—Mostly poorly sorted, clay-supported gravel in a sandy or silty matrix. May locally include clayey deposits in some subsidence troughs. Includes terraces up to about 12 ft above modern river level.
- Sheetwash deposits (Holocene and late Pleistocene)—Bubbly silty sand, sandy silt, and clayey silt deposited in ephemeral and intermittent stream valleys, on gentle hillslopes, and in basins.
- Younger terrace alluvium (late Pleistocene)—Mostly poorly sorted, clay-supported, locally bouldery pebble and cobble gravel in a sand and silt matrix. Deposited as glacial outwash. Underlies terraces 15 to 52 ft above modern stream level. May include fine-grained overbank deposits.
- Intermediate terrace alluvium (late Pleistocene)—Deposits texturally and depositionally similar to younger terrace alluvium (Qy). Underlies terraces 15 to 100 ft above modern streams.
- Oldest terrace alluvium (middle and early Pleistocene)—Deposits texturally and depositionally similar to younger terrace alluvium (Qy). Clasts moderately to highly well-sorted. A single small terrace remnant in the southeast corner of the quadrangle that is about 380 to 400 ft above the Frying Pan River.
- High-level gravel (early Pleistocene and/or late Tertiary)—Chiefly clay-supported, sandy silt, cobble and pebble gravel occurring on a subtle ridge line about 1 mi east of El Juel and about 1,300 to 1,500 ft above the Roaring Fork River. Clasts are moderately to very highly weathered.
- Sediments of Missouri Heights (early Quaternary and/or late Tertiary)—Locally derived gravel, sand, silt, and clay deposited in the Missouri Heights area in alluvial and colluvial environments. May include pediment deposits derived from and deposited on the sediments of Missouri Heights in areas between Spring Park Reservoir and Cattle Creek. Deposited in topographic depressions created by evaporite tectonism. Overlies Miocene basaltic rocks (Tb). Typically is less deformed than underlying rocks. Occurs about 1,000 to 1,650 ft above the Roaring Fork River.

MASS-WASTING DEPOSITS

- Recent landslide deposits (latest Holocene)—Includes a recently active landslide near the northeast corner of the map with very fresh morphological features. Heterogeneous unit consisting of unsorted, unstratified rock debris, sand, and silt.
- Colluvium (Holocene and late Pleistocene)—Ranges from unsorted, clay-supported, pebble to boulder gravel in a sandy silt matrix to matrix-supported gravelly, clayey, sandy silt. Usually coarser grained in upper reaches of colluvial slope and finer grained in distal areas.
- Talus (Holocene and late Pleistocene)—Angular, cobbly and bouldery rubble derived from outcrops of basalt, trachyandesite, and/or basaltic trachyandesite.
- Boulder-field deposits (Holocene and late Pleistocene)—Angular boulders and cobbles of basalt with little or no matrix in moderate to steep slopes. Commonly has an undulating surface suggestive of flowage as a rock glacier or related to periglacial processes.
- Landslide deposits (Holocene and Pleistocene)—Includes various types of landslide deposits. Consists of unsorted, unstratified gravel, sand, silt, clay, and rock debris.
- Older colluvium (Pleistocene)—Texturally similar to colluvium (Qc), but found on drainage divides, ridge lines, and dissected hillslopes.
- Older landslide deposits (Pleistocene)—Landslide deposits dissected by erosion and lacking distinctive landslide geomorphology. Similar in texture to landslide deposits (Ql).

ALLUVIAL AND MASS-WASTING DEPOSITS

- Younger debris-flow deposits (Holocene and late Pleistocene)—Poorly sorted to moderately well-sorted, matrix- and clay-supported deposits ranging from gravelly clayey silt to sandy, silty, cobbly, pebbly, and bouldery gravel. Fan heads tend to be bouldery, while distal fan areas are finer grained. Includes debris-flow, hyper-concentrated-flow, fluvial, and sheetwash deposits on active fans and in some drainage channels. Numerical subscripts indicate relative ages of younger debris-flow deposits in the southwest corner of the quadrangle. Deposits labeled Qdly are younger than and derived from deposits labeled Qdly.
- Alluvium and colluvium, undivided (Holocene)—Moderately well-sorted to well-sorted, stratified, interbedded sand, silt, pebbly sand, and sandy gravel to poorly sorted, unstratified or poorly stratified, clayey, silty sand, bouldery sand, sandy silt, and silty clay.
- Colluvium and sheetwash deposits, undivided (Holocene and late Pleistocene)—Consists of colluvium (Qc) on steeper slopes and sheetwash deposits (Qsw) on flatter slopes. Mapped where contacts between the two types of deposits are very gradational and difficult to locate. May locally include lacustrine deposits in large subsidence troughs.
- Older alluvium and colluvium, undivided (Pleistocene)—Deposits texturally and depositionally similar to alluvium and colluvium (Qac) that underlie terraces and hillslopes ranging from about 10 to 60 ft above the floor of tributary valleys.
- Older colluvium and sheetwash deposits, undivided (Pleistocene)—Deposits texturally and depositionally similar to colluvium and sheetwash (Qsw) that underlie surfaces 20 to 160 ft above adjacent stream beds.
- Older debris-flow deposits (Pleistocene)—Remnant of an inactive debris fan on a ridge line about 80 to 120 ft above the adjacent stream bed near the southeast corner of the quadrangle. Similar in texture and genesis to younger debris-flow deposits (Qdly).

SINTER DEPOSITS

- Tufa (Holocene and late Pleistocene)—Low-density, porous, calcium carbonate deposits precipitated from a mineral spring along the Basalt Mountain Fault immediately north of Cattle Creek.

UNDIFFERENTIATED SURFICIAL DEPOSITS

- Q Surficial deposits (Quaternary)—Shown only on cross section.

COLLAPSE DEPOSITS


- Collapse deposits (Quaternary and late Tertiary)—Heterogeneous deposits of moderately to highly deformed bedrock and overlying undeformed to moderately deformed surficial deposits within the Carbonadale Collapse Center. Locally includes large but displaced blocks of volcanic rocks. Formed in response to differential collapse resulting from dissolution and flow of underlying evaporite.

BEDROCK

- Trachyandesite of Spring Park (Pliocene)—Medium-gray basaltic flows from an eruptive center about 0.5 mi east of the dam for Spring Park Reservoir. Petrographically the unit is xenocrystic olivine basalt, while geochemically it is a basaltic trachyandesite. Groundmass predominantly plagioclase and pyroxene. Contains sparse phenocrysts of mainly olivine and rare plagioclase. Locally contains abundant xenocrysts of quartz, sandstone, and plagioclase. Included in compositional group 6b of Budahn and others (2002). ⁴⁰Ar/³⁹Ar of 2.96 ± 0.02 Ma (Kunk and others, 2002).
- Cinder deposits of Spring Park (Pliocene)—Red and red-brown scoriae, unconsolidated cinder deposits associated with an eroded eruptive center about 0.5 mi east of the dam for Spring Park Reservoir. Petrographically the rock is olivine basalt with locally abundant xenocrysts of quartz, sandstone, and plagioclase. Geochemically these rocks are basaltic trachyandesite and are included in compositional group 6b of Budahn and others (2002). ⁴⁰Ar/³⁹Ar of 2.96 ± 0.02 Ma (Kunk and others, 2002).
- Trachyandesite flows of Cattle Creek (Pliocene)—Medium- to dark-gray basaltic trachyandesite and trachybasalt flows along Cattle Creek between Shipes Drive and Sleepy Creek. Contains varying amounts of quartz and sandstone xenocrysts. Included in compositional group 6c of Budahn and others (2002). ⁴⁰Ar/³⁹Ar of 3.08 ± 0.02 Ma (Kunk and others, 2002).
- Trachyandesite cinder deposits of Cattle Creek (Pliocene)—Dark gray to black, scoriae, cinder deposit exposed in cinder quarries along Cattle Creek between Shipes Drive and Sleepy Creek. Petrographically this deposit is a xenocrystic, olivine basalt; geochemically it is basaltic trachyandesite. Included in compositional group 6c of Budahn and others (2002). ⁴⁰Ar/³⁹Ar of 3.01 ± 0.03 Ma (Kunk and others, 2002).
- Trachyandesite flows, undifferentiated (Pliocene)—Multiple flows of basaltic trachyandesite, trachybasalt, and trachybasalt. Contains varying amounts of quartz, sandstone, and plagioclase xenocrysts.
- Sediments of Basalt Mountain (Pliocene or Miocene)—Chiefly medium-red-brown, weakly indurated, pebble and cobble gravel in a sandy or silty matrix. Locally bouldery. Deposited over basalt flows on northern edge of Basalt Mountain shield volcano by ancestral Cattle Creek.
- Basaltic flows (Miocene)—Multiple flows of basalt, basaltic andesite, and basaltic trachyandesite. Petrographically most flows are olivine basalt; many are porphyritic. Groundmass predominantly plagioclase and pyroxene. Phenocrysts chiefly olivine and occasionally plagioclase. May contain rare xenocrysts or xenoliths of quartz and quartzite. Locally includes slightly indurated sediment. Included in compositional groups 1b, 1c, 2b, 4b, and 4c of Budahn and others (2002). ⁴⁰Ar/³⁹Ar age dates range from 7.75 ± 0.13 Ma to 10.84 ± 0.06 Ma.
- Sedimentary deposits (Miocene)—Mostly fluvial, clay-supported, silty, sandy pebble and cobble gravel but locally contains silty and sandy deposits of probable alluvial and/or colluvial origin. Locally slightly to moderately indurated.

- Manos Shale (Upper Cretaceous)—Light- to dark-gray, carbonaceous, silty to sandy shale and thin bentonite beds, gray limestone, and light- to medium-gray, grayish-yellow weathering, clayey sandstone. Includes the Fort Lyons Limestone Member, a thick-bedded, coarse-grained, gray limestone.
- Dakota Sandstone (Lower Cretaceous)—Light gray to tan, medium- to very coarse-grained, quartzose sandstone and conglomeratic sandstone interbedded with carbonaceous siltstone, sandstone, and shale. May include Burro Canyon Formation in southern part of quadrangle.
- Merion Formation (Upper Jurassic)—Pale-green, greenish-gray, and massive variegated siltstone and claystone, buff to tan sandstone, and gray limestone. A thick-bedded, coarse-grained, silty, tan and white weathering, medium- to dark-gray limestone at the base of the formation overlies the Fort Lyons Sandstone.
- Entrada Sandstone (Upper Jurassic)—Light gray, tan, and white, medium- to very fine-grained, well-sorted sandstone with large-scale crossbedding. Weakly to moderately indurated.
- Chinle Formation (Upper Triassic)—Thin, even-bedded, and structureless beds of dark reddish-brown, orangish-red, and purplish-red, calcareous siltstone and mudstone and scattered thin lenses of light purplish-red and gray limestone and limestone-pebble conglomerate. Locally includes a thin, basal conglomeratic sandstone.
- State Bridge Formation (Lower Triassic and Permian)—Reddish-orange, grayish-red, and pale-reddish-pink silty sandstone, clayey siltstone, arkosic sandstone, and conglomeratic sandstone. Includes lenses of sandy dolomite and limestone of the South Canyon Creek Dolomite Member that are up to 18 inches thick and occur about 200 ft above the base of the formation. Sandstone beds are well sorted, equigranular, and have rounded to sub-rounded sand grains with a high degree of sphericity.
- Chinle and State Bridge Formations, undivided (Triassic and Permian)
- Maroon Formation (Lower Permian? and Upper Pennsylvanian)—Red beds of sandstone, conglomerate, mudstone, siltstone, and shale and minor thin beds of gray limestone.
- Eagle Valley Formation (Middle Pennsylvanian)—Reddish-brown, gray, reddish-gray, and tan siltstone, shale, sandstone, gypsum, and carbonate rocks which are gradational between and intertonguing with the Maroon Formation and Eagle Valley Evaporite.
- Maroon and Eagle Valley Formations, undivided (Lower Permian and Upper and Middle Pennsylvanian)
- Eagle Valley Evaporite (Middle Pennsylvanian)—Evaporitic sequence of gypsum, anhydrite, and halite interbedded with mudstone, fine-grained sandstone, thin carbonate beds, and black shale. Commonly intensely folded, faulted, and ductily deformed.

CORRELATION OF MAP UNITS



ACKNOWLEDGEMENTS

This geologic map was funded in part by the U.S. Geological Survey National Geologic Mapping Program and by the State of Colorado General Funds and Colorado Department of Natural Resources Severance Tax Operational Funds.